AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims

passion flower.

Claim 1 (Currently Amended): A pharmaceutical composition comprising:

an effective amount of a β (1-3) β (1-4) glucan composition consisting essentially of at least

about 75% β (1-3) β (1-4) glucan having a particle size of less than 0.2 μm, less than 10% ash impurities, less than 10% protein impurities and less than 5% lipid impurities, and

an effective amount of a botanical extract, or a pharmaceutically active agent.

Claim 2 (Original): The pharmaceutical composition according to claim 1, wherein the composition comprises the botanical extract, and wherein the botanical extract is an extract of Guarana, Ginkgo biloba, Kola nut, Goldenseal, Golo Kola, Schizandra, Elderberry, St. John's Wort, Valerian and Ephedra, black tea, white tea, java tea, garlic oil, fiber, green tea, lemon oil, mace, licorice, onion oil, orange oil, rosemary, milk thistle, Echinacea, Siberian ginseng or Panax ginseng, lemon balm, Kava kava, matte, bilberry, soy, grapefruit, seaweed, hawthorn, lime blossom, sage, clove, basil, curcumin, taurine, wild oat herb, oat grain, dandelion, gentian, aloe vera, hops, cinnamon, peppermint, grape, chamomile, fennel, marshmallow, ginger, slippery elm, cardamon, coriander, anise, thyme, rehmannia, eucalyotus, menthol, schisandra, withania, cowslip, lycium, or

Claim 3 (Original): The pharmaceutical composition according to claim 2, wherein the botanical extract is an extract of oat grain.

Application Serial No. 10/554,290 Amendment dated 3 March 2008

Reply to Office Action mailed 3 December 2007

Claim 4 (Withdrawn): The pharmaceutical composition of claim 3, wherein the botanical

extract comprises avenanthramide.

Claim 5 (Withdrawn): The pharmaceutical composition according to claim 1, wherein the

composition comprises the pharmaceutically active agent, and wherein the pharmaceutically active agent is selected from the group consisting of beta-sitosterol, caffeine, cafestol, D-limonene,

kabweol, nomilin, oltipraz, sulphoraphane, tangeretin, folic acid, and menthol.

Claim 6 (Withdrawn): The pharmaceutical composition according to claim 1, wherein the

composition comprises the pharmaceutically active agent, and wherein the pharmaceutically active agent is selected from the group consisting of an antihistamine, a decongestant, a corticosteroid, a

non-steroidal anti-inflammatory drug, a bronchodilator, a vasodilator, such as nitroglycerin, and a

local anaesthetic.

Claim 7 (Withdrawn): The pharmaceutical composition of claim 6, wherein the vasodilator

is nitroglycerin.

Claim 8 (Original): The pharmaceutical composition according to claim 1, wherein the β

(1-3) β (1-4) glucan is derived from a cereal grain or a part of the cereal grain.

Claim 9 (Original): The pharmaceutical composition according to claim 8, wherein the

cereal is selected from the group consisting of a cultivar of barley, a cultivar of oat, a cultivar of

wheat, a cultivar of rye, a cultivar of sorghum, a cultivar of millet, a cultivar of com, and a mixture thereof.

Claim 10 (Canceled).

Page 7 of 17

Claim 11 (Currently Amended): The pharmaceutical composition according to claim ± 0.1 , wherein the β (1-3) β (1-4) glucan composition has a purity of at least about 92%, and contains less than 3.5% ash impurities, less than 3.5% protein impurities, and less than 1% lipid impurities.

Claim 12 (Currently Amended): The pharmaceutical composition according to claim $^{+0}$ 1, herein the cereal β -glucan composition has a clarity value of from about 5 to about 100 NTU

Claim 13 (Currently Amended): The pharmaceutical composition according to claim 1, wherein the β (1-3) β (1-4) glucan <u>composition</u> is produced according to a method of isolating a β (1-3) β (1-4) glucan <u>composition</u> from a milled cereal grain or a milled part of the cereal grain, the method comprising:

- (i) extracting the milled cereal grain or the milled part of the cereal grain with an alkaline solution <u>having a value of pH of between 9 to 10 for a period of time of about 15 to about 45 minutes</u> to produce an extract containing at least about 0.4 weight percent β (1-3) β (1-4) glucan;
- (ii) removing insoluble material, and removing particulate material having a particle size of greater than about $0.2~\mu m$ from said extract to produce a purified extract comprising β (1-3) β (1-4) glucan having a particle size of less than $0.2~\mu m$ wherein the step of removing particulate material comprises using filtering out material having a particle size of greater than about $0.2~\mu m$ from said extract by microfiltration to produce the purified extract comprising β (1-3) β (1-4) glucan having a particle size of less than $0.2~\mu m$ as a filtrate;
- (iii) adding from about 10% to about $\frac{25\% (w/w)}{20\% (vol/vol)}$ of a C_1 - C_4 alcohol to the purified extract to precipitate the β (1-3) β (1-4) glucan composition, and
 - (iv) isolating the β (1-3) β (1-4) glucan <u>composition</u>.

Claim 14 (Currently Amended): The pharmaceutical composition according to claim 13, wherein in said step of adding (step iii) in said method, about 10% to about 20% (w/w) of an the alcohol is selected from the group consisting of methanol, ethanol and isopropanol, is used to precipitate the B (1-3) B (1-4) glucan from said purified extract.

Claim 15 (Currently Amended): The pharmaceutical composition according to claim 14, wherein the alcohol about 10% to about 20% (w/w) of is ethanol is used to precipitate the β (1-3) β (1-4) glucan from said purified extract.

Claim 16 (Currently Amended): The pharmaceutical composition according to claim 13, wherein said step of removing particulate material in said method <u>further</u> comprises <u>the following</u> steps prior to the step of microfiltration:

one, or more than one step of adding a flocculant, a coagulant or both α the flocculant and α the coagulant to said extract to coagulate particulate material having a particle size of greater than about 0.2 μ m, and removing coagulated material from said extract, and

digesting starch material in said extract, and

filtering out particulate material having a particle size of greater than about 0.2 mm from said extract to produce a purified extract.

Claim 17 (Original): The pharmaceutical composition according to claim 16, wherein in said step of digesting in said method, said starch material is digested with an enzyme.

Claim 18 (Original): The pharmaceutical composition according to claim 17, wherein prior to digesting said starch material, said alkaline solution is neutralized.

Claim 19 (Original): The pharmaceutical composition according to claim 18, wherein following the digestion of said starch material in said method, said enzyme is inactivated.

Claim 20 (Original): The pharmaceutical composition according to claim 19, wherein said

enzyme is inactivated by acidifying the neutralized solution.

Claim 21 (Original): The pharmaceutical composition according to claim 17, wherein said

enzyme is an amylase.

Claim 22 (Original): The pharmaceutical composition according to claim 21, wherein said

amylase does not require a calcium cofactor.

Claim 23 (Original): The pharmaceutical composition according to claim 13, wherein the

cereal is selected from the group consisting of a cultivar of barley, a cultivar of oat, a cultivar of wheat, a cultivar of rye, a cultivar of sorghum, a cultivar of millet, a cultivar of corn, and a mixture

thereof.

Claim 24 (Currently Amended): The pharmaceutical composition according to claim 13,

wherein the pH of the alkaline solution used in said method is from about 9 9.25 to about 10 9.75.

Claim 25 (Canceled).

Claim 26 (Original): The pharmaceutical composition according to claim 13, wherein said

step of adding (step iii) in said method is conducted at a temperature of from about 1°C to about

10°C.

Page 10 of 17

Claim 27 (Currently Amended): The pharmaceutical composition according to claim 13, wherein said method further comprises one, or more than one step of dissolving the isolated β (1-3) β (1-4) glucan in an aqueous solution, precipitating the β (1-3) β (1-4) glucan by adding about 10% to about $\frac{25\% (w/w)}{20\% (vol/vol)}$ of the C_1 - C_4 alcohol to the aqueous solution, and isolating the β (1-3) β (1-4) glucan.

Claim 28 (New): A pharmaceutical composition consisting essentially of:

- a β (1-3) β (1-4) glucan composition consisting essentially of at least about 75% β (1-3) β (1-4) glucan having a particle size of less than 0.2 μ m, less than 10% ash impurities, less than 10% protein impurities and less than 5% lipid impurities.
 - a botanical extract, or a pharmaceutically active agent, and
 - a pharmaceutically acceptable diluent or carrier.

Claim 29 (New): The pharmaceutical composition according to claim 28, wherein the β (1-3) β (1-4) glucan composition has a purity of at least about 92%, and contains less than 3.5% ash impurities, less than 3.5% protein impurities, and less than 1% lipid impurities.

Claim 30 (New): A pharmaceutical composition consisting of:

- a β (1-3) β (1-4) glucan composition consisting essentially of at least about 75% β (1-3) β (1-4) glucan having a particle size of less than 0.2 μ m, less than 10% ash impurities, less than 10% protein impurities and less than 5% lipid impurities.
 - a botanical extract, or a pharmaceutically active agent, and
 - a pharmaceutically acceptable diluent or carrier.

Application Serial No. 10/554,290 Amendment dated 3 March 2008 Reply to Office Action mailed 3 December 2007

Claim 31 (New): The pharmaceutical composition according to claim 30, wherein the β (1-3) β (1-4) glucan composition has a purity of at least about 92%, and contains less than 3.5% ash impurities, less than 3.5% protein impurities, and less than 1% lipid impurities.